

Vertex Degree of Random Intersection Graph

by

Bhupendra Gupta ¹

Faculty of Engineering and Sciences,
Indian Institute of Information Technology (DM)-Jabalpur, India.

Abstract

A random intersection graph is constructed by independently assigning a subset of a given set of objects W , to each vertex of the vertex set V of a simple graph G . There is an edge between two vertices of V , iff their respective subsets(in W ,) have at least one common element. The strong threshold for the connectivity between any two arbitrary vertices of vertex set V , is derived. Also we determine the almost sure probability bounds for the vertex degree of a typical vertex of graph G .

Keywords: Random intersection graphs, Vertex degree.

AMS 2000 subject classifications: 05C80, 91D30.

¹Corresponding Author. email:gupta.bhupendra@gmail.com, bhupen@iiitdm.in

1 Introduction

The development in this paper is in continuation of Dudley Stark [2], in which distribution of the degree of a typical vertex is given. Dudley studied the model given in [3] defined as:

Let us consider a set V with n vertices and another set of objects W with m objects. Define a bipartite graph $G^*(n, m, p)$ with independent vertex sets V and W . Edges between $v \in V$ and $w \in W$ exists independently with probability p . The random intersection graph $G(n, m, p)$ derived from $G^*(n, m, p)$ is defined on the vertex set V with vertices $v_1, v_2 \in V$ are adjacent if and only if there exists some $w \in W$ such that both v_1 and v_2 are adjacent to w in $G^*(n, m, p)$. Also define W_v be a random subset of W such that each element of W_v is adjacent to $v \in V$. Any two vertices $v_1, v_2 \in V$ are adjacent if and only if $W_{v_1} \cap W_{v_2} \neq \phi$, and edge set $E(G)$ is define as

$$E(G) = \{\{v_i, v_j\} : v_i, v_j \in V, W_{v_i} \cap W_{v_j} \neq \phi\}.$$

2 Definitions and Supporting Results

Lemma 2.1 *Let X be a random variable having binomial distribution with parameters n and $p > 0$, i.e., $X \sim Bi(n, p)$. Then*

$$P[X \geq k] \leq \left(\frac{np}{k}\right)^k \exp(k - np), \quad k \geq np, \quad (2.1)$$

and

$$P[X \leq k] \leq \left(\frac{np}{k}\right)^k \exp(k - np), \quad 0 < k \leq np. \quad (2.2)$$

For convenience, the bounds (2.1) and (2.2) can be expressed as

$$\exp\left(npH\left(\frac{np}{k}\right)\right),$$

where

$$H(t) = \frac{1}{t} \log t + \frac{1}{t} - 1, \quad 0 < t < \infty,$$

and $H(\infty) = -1$. Note that $H(t) < 0, \forall t \in (0, \infty] \setminus \{1\}$; H is increasing on $(0, 1)$ and decreasing on $(1, \infty)$. (See Lemma 1.2 on page 25, Penrose [4].) $\square$

Let $X = X(n, m, p)$ be the number of vertices of $V - \{v\}$ adjacent in $G(n, m, p)$ to a vertex $v \in V$, i.e., X be the vertex degree of a vertex $v \in V$ in $G(n, m, p)$. Then X follows $\text{Bi}(n-1, q_n)$, where q_n is the probability of $v_1, v_2 \in V$ are adjacent.

Proposition 2.2 *Let $v_i, v_j \in V, i \neq j$ and $i, j = 1, 2, \dots, n$. Then v_i and v_j are adjacent with probability q_n , such that*

$$q_n \sim mp^2,$$

for sufficiently small p .

Proof. Consider

$$\begin{aligned} q_n &= P[v_i, v_j \text{ are adjacent.}] \\ &= P[W_{v_i} \cap W_{v_j} \neq \emptyset] \\ &= P[|W_{v_i} \cap W_{v_j}| \geq 1] \\ &= 1 - P[w \notin (W_{v_i} \cap W_{v_j})]^m \\ &= 1 - [1 - P[w \in (W_{v_i} \cap W_{v_j})]]^m \\ &= 1 - [1 - P[w \in W_{v_i}, w \in W_{v_j}]]^m \\ &= 1 - [1 - p^2]^m \end{aligned}$$

Using the Taylor's series expansion up to second term, we get

$$q_n = mp^2 - \zeta_m, \tag{2.3}$$

where, $\zeta_m = \frac{m(m-1)}{2!}(1-c)^2p^4$, and $c \in (0, p^2)$.

Now if we take p is sufficiently small. Then

$$q_n \sim mp^2. \tag{2.4}$$

This completes the proof. $\square$

Definition 1 *Let graphs A and B share the same vertices and the edge set of A is a subset of the edge set of B , we write $A \leq B$. Also let Θ be a property of a random graphs such that if $A \leq B$ and $A \in \Theta$, then $B \in \Theta$. (Here $A \in \Theta$ is used to denote that graph A has property Θ .) Then Θ is called an upwards-closed property. If $B \in \Theta$ implies $A \in \Theta$, then Θ is said to be a downwards-closed property. If property Θ is upwards-closed or downwards-closed, then Θ is called monotone property. $\square$*

Fix a monotone property Θ . For any two functions $\delta, \gamma : Z^+ \rightarrow R^+$, we write $\delta \ll \gamma$ (resp. $\delta \gg \gamma$) if $\delta(n)/\gamma(n) \rightarrow 0$, (resp. $\gamma(n)/\delta(n) \rightarrow 0$) as $n \rightarrow \infty$. We will write δ for $\delta(n)$. Let $G_n(r)$, be any random graph on n vertices.

Definition 2 *A function $\delta_\Theta : Z^+ \rightarrow R^+$ is a weak threshold function for Θ if the following is true for every function $\delta : Z^+ \rightarrow R^+$,*

- *if $\delta(n) \ll \delta_\Theta(n)$, then $P[G_n(\delta) \in \Theta] = 1 - o(1)$, and*
- *if $\delta(n) \gg \delta_\Theta(n)$ then $P[G_n(\delta) \in \Theta] = o(1)$. $\square$*

A function $\delta_\Theta : Z^+ \rightarrow R^+$ is a strong threshold function for Θ if the following is true for every fixed $\epsilon > 0$,

- *if $P[G_n(\delta_\Theta - \epsilon) \in \Theta] = 1 - o(1)$, and*
- *if $P[G_n(\delta_\Theta + \epsilon) \in \Theta] = o(1)$. $\square$*

3 Main Results

We now derive the threshold probability for the connectivity of graph $G(n, m, p)$.

Let Θ be the connectivity of graph $G(n, m, p)$ and define the probability that a vertex $v \in V$ is connected to $w \in W$, as follows

$$p := p(\alpha) = \frac{1}{(mn^\alpha)^{1/2}} .$$

Theorem 3.1 *Let $G(n, m, p)$ be the random intersection graph with $p(\alpha) = \frac{1}{(mn^\alpha)^{1/2}}$ and $v_i, v_j \in V(G)$ for $i \neq j = 1, 2, \dots, n$. Then $p(2) = \frac{1}{m^{1/2}n}$, is a strong threshold for the random intersection graph $G(n, m, p)$, i.e.,*

$$\begin{aligned} P[G(n, m, p(2 - \epsilon)) \in \Theta] &= 1 - o(1), \\ P[G(n, m, p(2 + \epsilon)) \in \Theta] &= o(1). \end{aligned} \tag{3.5}$$

Proof. Let $v_i, v_j \in V(G)$ for $i \neq j, i, j = 1, 2, \dots, n$. Then from (2.3), we have

$$\begin{aligned} P[v_i, v_j \text{ are adjacent}] &= q_n \\ &\leq mp^2. \end{aligned} \tag{3.6}$$

Since we have $p(\alpha) = \frac{1}{(mn^\alpha)^{1/2}}$. Then

$$P[v_i, v_j \text{ are adjacent}] \leq n^{-\frac{\alpha}{2}}. \tag{3.7}$$

Hence, for $\alpha \leq 2$ the above probability (3.7) is not summable, i.e.,

$$\sum_{n=0}^{\infty} P[v_i, v_j \text{ are adjacent}] = \infty.$$

Since the adjacency between any pair of vertices is independent from the adjacency between any other pair of vertices. Then by the Borel-Cantelli Lemma, we have

$$P[v_i, v_j \text{ are adjacent}, \quad i.o.] = 1.$$

Therefore, for $\alpha \leq 2$, any pair of vertices $v_i, v_j \in V, i \neq j, i, j = 1, 2, \dots, n$, are adjacent almost surely. Hence,

$$P[G(n, m, p(2 - \epsilon)) \in \Theta] = 1 - o(1).$$

For $\alpha > 2$ the above probability is (3.7) summable, i.e.,

$$\sum_{n=0}^{\infty} P[v_i, v_j \text{ are adjacent}] < \infty.$$

Then by the Borel-Cantelli's Lemma, we have

$$P[v_i, v_j \text{ are adjacent, } i.o.] = 0.$$

Therefore, for $\alpha > 2$, any pair of vertices $v_i, v_j \in V$ are adjacent only finitely many times. Hence,

$$P[G(n, m, p(2 + \epsilon)) \in \Theta] = o(1).$$

This completes the proof. □

We now state strong law results for vertex degree of a typical vertex.

Let $X = X(n, m, p)$ be the number of vertices of $V - \{v\}$ adjacent in $G(n, m, p)$ to a vertex $v \in V$, $i, j = 1, 2, \dots, n$, $i \neq j$ i.e., X be the degree of vertex $v \in V$ in $G(n, m, p)$.

Theorem 3.2 *Let $G(n, m, p)$ be the random intersection graph with $p = (mn^\alpha)^{-1/2}$, where $0 < \alpha < 1$. Also let X be the degree of a typical vertex $v \in V$ in $G(n, m, p)$. Then*

$$\limsup_{n \rightarrow \infty} \frac{X}{n^\delta} \geq a(c), \quad a.s., \quad (3.8)$$

where $\delta = 1 - \alpha$ and $a(c)$ is the root in $[1, \infty)$ of

$$a \log a - a + 1 = c, \quad (3.9)$$

with $a(\infty) = 1$.

Proof. We know that the vertex degree of a vertex in graph $G(n, m, p)$ is distributed according to $\text{Bi}(n - 1, q_n)$. Hence by Lemma 2.1, we have

$$P[X \leq K] \leq \exp \left((n - 1)q_n H \left(\frac{(n - 1)q_n}{K} \right) \right), \quad K \leq nq_n. \quad (3.10)$$

From Proposition 2.2, we have for sufficiently small p , we have $q_n \sim mp^2$. Then

$$\begin{aligned} P[X \leq K] &\leq \exp \left((n-1)mp^2 H \left(\frac{(n-1)mp^2}{K} \right) \right) \\ &\sim \exp \left(n^{1-\alpha} H \left(\frac{n^{1-\alpha}}{K} \right) \right) \\ &= \exp \left(-cn^\delta \right), \end{aligned} \tag{3.11}$$

where $H((n/mK^2)^{1/2}) = -c$ and $\delta = 1 - \alpha$. The above expression is summable if $K = a(c)n^\delta$, where $a(c)$ is increasing in $[1, \infty)$ and defined as in (3.9). Then by the Borel-Cantelli Lemma we have

$$P \left[X \leq a(c)n^\delta, \quad i.o. \right] = 0. \tag{3.12}$$

This implies that

$$\limsup_{n \rightarrow \infty} \frac{X}{n^\delta} \geq a(c), \quad a.s. \tag{3.13}$$

This completes the proof. $\square$

Theorem 3.3 *Let $G(n, m, p)$ be the random intersection graph with $p = (mn^\alpha)^{-1/2}$, where $0 < \alpha < 1$. Also let X be the degree of a typical vertex $v \in V$ in $G(n, m, p)$. Then*

$$\liminf_{n \rightarrow \infty} \frac{X}{n^\delta} \leq a(c), \quad a.s., \tag{3.14}$$

where $\delta = 1 - \alpha$ and $a(c)$ is the root in $(0, 1)$ of

$$a \log a - a + 1 = c, \tag{3.15}$$

with $a(\infty) = 1$.

Proof. We know that the vertex degree of a vertex in graph $G(n, m, p)$ is distributed according to $\text{Bi}(n-1, q_n)$. Hence by Lemma 2.1 we have

$$P[X \geq K] \leq \exp \left((n-1)q_n H \left(\frac{(n-1)q_n}{K} \right) \right), \quad K \geq nq. \tag{3.16}$$

From Proposition 2.2, we have for sufficiently small p , we have $q_n \sim mp^2$. Then

$$\begin{aligned} P[X \geq K] &\leq \exp \left((n-1)mp^2 H \left(\frac{(n-1)mp^2}{K} \right) \right) \\ &\sim \exp \left(n^{1-\alpha} H \left(\frac{n^{1-\alpha}}{K} \right) \right) \\ &= \exp \left(-cn^\delta \right), \end{aligned} \tag{3.17}$$

where $H((n/mK^2)^{1/2}) = -c$ and $\delta = 1 - \alpha$. The above expression is summable if $K = a(c)n^\delta$, where $a(c)$ is decreasing in $(0, 1)$ and defined as in (3.15). Then by the Borel-Cantelli Lemma, we have

$$P \left[X \geq a(c)n^\delta, \quad i.o. \right] = 0. \tag{3.18}$$

This implies that

$$\liminf_{n \rightarrow \infty} \frac{X}{n^\delta} \leq a(c), \quad a.s. \tag{3.19}$$

This completes the proof. $\square$

References

- [1] Y.S.Chow, H.Teicher. Probabilty Theory, thired edition, Springer Text in Statistics (2004).
- [2] D. Stark. The vertex degree distribution of random intersection graphs, Random Structures and Algorithms, **Vol. 24**(3): 249 - 258 , (2004).
- [3] M. Karonski, E.R.Scheinerman and K.B.Singer-Cohen. On random intersection graphs: the subgraph problem, Combinatorics, Probability and Computing, **Vol. 8**: 131-159 (1999).
- [4] Penrose, M.D.(2003), Random Geometric Graphs, second edition, Oxford University Press.